

Pathology Job Interview Questions And Answers



Interview Questions Answers

<https://interviewquestionsanswers.org/>

About Interview Questions Answers

Interview Questions Answers . ORG is an interview preparation guide of thousands of Job Interview Questions And Answers, Job Interviews are always stressful even for job seekers who have gone on countless interviews. The best way to reduce the stress is to be prepared for your job interview. Take the time to review the standard interview questions you will most likely be asked. These interview questions and answers on Pathology will help you strengthen your technical skills, prepare for the interviews and quickly revise the concepts.

If you find any **question or answer** is incorrect or incomplete then you can **submit your question or answer** directly with out any registration or login at our website. You just need to visit [Pathology Interview Questions And Answers](#) to add your answer click on the *Submit Your Answer* links on the website; with each question to post your answer, if you want to ask any question then you will have a link *Submit Your Question*; that's will add your question in Pathology category. To ensure quality, each submission is checked by our team, before it becomes live. This [Pathology Interview preparation PDF](#) was generated at **Wednesday 29th November, 2023**

You can follow us on FaceBook for latest Jobs, Updates and other interviews material.
www.facebook.com/InterviewQuestionsAnswers.Org

Follow us on Twitter for latest Jobs and interview preparation guides.
<https://twitter.com/InterviewQA>

If you need any further assistance or have queries regarding this document or its material or any of other inquiry, please do not hesitate to contact us.

Best Of Luck.

Interview Questions Answers.ORG Team
<https://InterviewQuestionsAnswers.ORG/Support@InterviewQuestionsAnswers.ORG>



Pathology Interview Questions And Answers Guide.

Question - 1:

Artificial intelligence and data-mining systems often use "training data sets" and "test data sets." Define these terms and describe briefly how these data sets are used.

Ans:

Training data sets are given to systems initially to teach them to make correct responses. Test data sets are equivalent to the training sets but contain separate data and are used to verify the performance of the systems.

[View All Answers](#)

Question - 2:

Rule-based systems underlie most clinical event monitors (programs that detect important clinical events and notify appropriate medical personnel). Often these systems work in conjunction with data from the clinical pathology LIS. What aspects of clinical pathology make a rule-based system a reasonable approach?

Ans:

Clinical laboratory databases consist of many discrete test results that have known reference ranges and critical values. Well-established patterns of these results exist that is known as related to important clinical conditions. Writing rules that detect and alert to these patterns is straightforward.

[View All Answers](#)

Question - 3:

You are working with an intensive care unit (ICU) attending physician on a project to see if you can predict readmission for patients with pancreatitis. You have access to a large database of ICU data (such as cardiac catheter values, vital signs, and respiratory parameters), as well as all of the data that can be gleaned from the LIS. There are approximately 800 measurements of various types for each of 4000 patients. You do not really have any specific ideas about what values would be most predictive; in fact, you think it is likely that the predictors are highly complex combinations of factors. Which of the 3 types of artificial intelligence systems would be most appropriate for this problem, and why?

Ans:

A neural network is most appropriate, because there is no prior knowledge to allow selection of predictors, the relative weighting of predictors is unknown, a large data set of many discrete potential predictors is available, combinations of predictors may provide better discrimination than individual predictors, and the desired classification is binary (readmission likely or unlikely).

[View All Answers](#)

Question - 4:

What is the Arden syntax?

Ans:

The Arden syntax is a standard language and format for representing the medical knowledge and algorithms required for making medical decisions. It is used in medical decision support systems.

[View All Answers](#)

Question - 5:

How are neural networks different than Bayesian belief networks along the following dimensions:

- (1) inspect ability of knowledge,
- (2) need for probabilities acquired from "domain" experts,
- (3) need for data to train the system, and
- (4) ability of the system to make classifications based on input data. (Note: You may find it helpful to make a 2 \times 4 table and include a short phrase or two in each cell.)

Ans:

Bayesian belief networks are inspectable, known probabilities are required, training data are not needed, and they can classify into multiple categories.

Neural networks are not inspectable, they do not need domain expertise or known probabilities, training data are required, and they are best for a binary classification ("yes" or "no").

[View All Answers](#)

Question - 6:



Most of the artificial intelligence systems we discussed rely on some kind of knowledge representation, with the notable exception of neural networks. Where is the "knowledge" in a neural network stored?

Ans:

In the weightings between: the nodes or "neurons."

[View All Answers](#)

Question - 7:

What is an "entity-relationship" diagram useful for, state briefly?

Ans:

An entity-relationship diagram is a way of illustrating the structure of a relational database in a simple format. It displays the primary "entities" (tables) in the database and the relationships that exist between the data elements in the tables. It is useful as a basis for discussion during database design and in describing existing databases.

[View All Answers](#)

Question - 8:

Describe the main difference between the hypothesis-testing and hypothesis-generating approaches to data mining.

Ans:

In hypothesis testing, data mining is used to determine whether and under what conditions a proposed pattern exists in a large data set. In hypothesis generation, data mining is used to discover patterns in the data without prior knowledge of what kinds of patterns might exist.

[View All Answers](#)

Question - 9:

What are process measures in outcomes research and why are they sometimes used in place of actual outcomes data?

Ans:

A process measure is a piece of data that is closely related to an outcome, but is easier to measure or more available than the actual outcome data. Thus, it is convenient to use as a surrogate measure for the outcome. For example, the effect of diabetes health education program, the number of eye examinations and regular evaluation of glycosylated hemoglobin (i.e., good practices) rather than assessing the actual long-term health of the diabetics.

[View All Answers](#)

Question - 10:

What advantage does a pathologist have over investigators in most other fields in carrying out outcomes or data-mining studies?

Ans:

Some of the most important and useful data in clinical data mining are derived from pathology services (anatomic pathology diagnoses and laboratory test results). In most places, pathologists manage the systems that contain these key data.

[View All Answers](#)

Question - 11:

Define "association rules" and describe their use in exploratory data mining.

Ans:

Association rules express the likelihood of co occurrence of features or events in records in a database (e.g. if a patient has characteristics A and B, he or she has an 80% chance of having characteristic C). Data-mining software can automatically identify associations in large data sets. Although many associations are trivial, some indicate causative or "common cause" relationships. Changing associations over time may also provide useful information.

[View All Answers](#)

Question - 12:

Outcomes research is often limited in the conclusions that can be drawn because of limitations in the data sources used for the studies. What are the most common data sources and what are their main limitations?

Ans:

The most common data sources include large local or regional administrative databases from hospitals, insurers, or government agencies. These databases contain very limited clinical information (usually ICD-9 codes), and thus it is difficult to meaningfully stratify patients by the severity of their illness, particular symptoms or test result characteristics, or the details of their therapy.

[View All Answers](#)

Question - 13:

Amyloid reacts with the Prussian blue stains.

Ans:

False

[View All Answers](#)

Question - 14:

Amyloid reacts with the alcian blue stains.

Ans:

False



[View All Answers](#)

Question - 15:

Amyloid reacts with the methyl violet stains.

Ans:

True

[View All Answers](#)

Question - 16:

Amyloid reacts with the thioflavine -T stains.

Ans:

True

[View All Answers](#)

Question - 17:

Amyloid reacts with the Congo red stains.

Ans:

True

[View All Answers](#)

Question - 18:

Necrosis is a feature in leprosy.

Ans:

False

[View All Answers](#)

Question - 19:

Necrosis is a feature in tuberculosis.

Ans:

True

[View All Answers](#)

Question - 20:

Necrosis is a feature in sarcoidosis.

Ans:

False

[View All Answers](#)

Question - 21:

Necrosis is a feature in syphilis.

Ans:

True

[View All Answers](#)

Question - 22:

Necrosis is a feature in actinomycosis.

Ans:

True

[View All Answers](#)

Question - 23:

Anaphylaxis causes urticaria.

Ans:

True

[View All Answers](#)

Question - 24:

Anaphylaxis is produced by Ig E antibody.

Ans:



True

[View All Answers](#)

Question - 25:

Anaphylaxis causes degranulation of basophils and mast cells.

Ans:

True

[View All Answers](#)

Question - 26:

Anaphylaxis causes eosinophilia.

Ans:

True

[View All Answers](#)

Question - 27:

Anaphylaxis occurs in 24 hours after the initial stimulus.

Ans:

False

[View All Answers](#)

Question - 28:

Collagens production is impaired in vitamin C deficiency.

Ans:

True

[View All Answers](#)

Question - 29:

Collagens synthesis is impaired in the presence of glucocorticoids.

Ans:

True

[View All Answers](#)

Question - 30:

The total amount of collagen in a wound reaches normal level within 4 weeks.

Ans:

False

[View All Answers](#)

Question - 31:

Collagens are secreted into the extracellular space as tropocollagen.

Ans:

True

[View All Answers](#)

Question - 32:

Collagens are produced in the endoplasmic reticulum of the fibroblasts.

Ans:

True

[View All Answers](#)

Question - 33:

Phagocytosis microorganisms are killed by IgG.

Ans:

False

[View All Answers](#)

Question - 34:

Phagocytosis microorganisms are killed by complement system.

Ans:



False

[View All Answers](#)

Question - 35:

Phagocytosis microorganisms are killed by lymphokines.

Ans:

False

[View All Answers](#)

Question - 36:

Phagocytosis microorganisms are killed by lysozyme.

Ans:

True

[View All Answers](#)

Question - 37:

Phagocytosis microorganisms are killed by hydrogen peroxide.

Ans:

True

[View All Answers](#)

Question - 38:

Malignant melanoma is found exclusively in the skin.

Ans:

False

[View All Answers](#)

Question - 39:

In malignant melanoma, staging is according to the size of the tumor.

Ans:

False

[View All Answers](#)

Question - 40:

In malignant melanoma, malignant transformation in common moles is about 1:10,000.

Ans:

False

[View All Answers](#)

Question - 41:

In malignant melanoma, amelanotic type is more aggressive than pigmented type.

Ans:

True

[View All Answers](#)

Question - 42:

Malignant melanoma is commoner in females than males.

Ans:

False

[View All Answers](#)

Question - 43:

An internal carotid aneurysm in the cavernous sinus can give rise to ipsilateral miosis.

Ans:

True

[View All Answers](#)

Question - 44:

Berry aneurysm is associated with hypercholesterolaemia.

Ans:



False

[View All Answers](#)

Question - 45:

Hypertension is associated with formation of true aneurysm.

Ans:

True

[View All Answers](#)

Question - 46:

Dissecting aneurysm results from destruction of the vessel media layer.

Ans:

True

[View All Answers](#)

Question - 47:

A true aneurysm is covered by all three layers of the vessel.

Ans:

True

[View All Answers](#)

Question - 48:

In Dermoid, cyst rupture of the cyst is associated with foreign body type inflammation.

Ans:

True

[View All Answers](#)

Question - 49:

In most cases dermoid cyst are congenital.

Ans:

True

[View All Answers](#)

Question - 50:

In Dermoid, cyst hair may be found in the cyst.

Ans:

True

[View All Answers](#)

Question - 51:

Dermoid cyst is lined by squamous epithelium.

Ans:

True

[View All Answers](#)

Question - 52:

If dermoid cyst found in the orbital region is usually located in the medial angle of the eye.

Ans:

False

[View All Answers](#)

Question - 53:

Sebaceous cyst contains mucopolysaccharides.

Ans:

False

[View All Answers](#)

Question - 54:

Sebaceous cyst is firmly adheres to the skin.

Ans:



False

[View All Answers](#)

Question - 55:

In Sebaceous cyst, scalp is a common location.

Ans:

True

[View All Answers](#)

Question - 56:

In Sebaceous cyst, central punctum is a common feature

Ans:

True

[View All Answers](#)

Question - 57:

Sebaceous cyst arises in the dermis.

Ans:

True

[View All Answers](#)

Question - 58:

In Squamous cell carcinoma, the tumours typically have everted edges.

Ans:

True

[View All Answers](#)

Question - 59:

In Squamous cell carcinoma, metastasis is usually to the regional lymph nodes.

Ans:

True

[View All Answers](#)

Question - 60:

Squamous cell carcinoma is the most common skin tumor seen in transplant patient.

Ans:

True

[View All Answers](#)

Question - 61:

Squamous cell carcinoma only occurs in the skin.

Ans:

False

[View All Answers](#)

Question - 62:

Squamous cell carcinoma is a commoner malignant skin tumor than basal cell carcinoma.

Ans:

False

[View All Answers](#)

Question - 63:

In corneal transplantation in the UK, HLA matching is routinely carried out before use.

Ans:

False

[View All Answers](#)

Question - 64:

In corneal transplantation in the UK, specular microscopy is used to count the number of endothelial cells.

Ans:



True

[View All Answers](#)

Question - 65:

In corneal transplantation in the UK, donor eyes older than 65 years of age are rejected.

Ans:

False

[View All Answers](#)

Question - 66:

In corneal transplantation in the UK, the cornea is kept at sub-zero temperature in the bank.

Ans:

False

[View All Answers](#)

Question - 67:

In corneal transplantation in the UK, whole eye is kept in the culture medium in the eye bank.

Ans:

False

[View All Answers](#)

Question - 68:

Berry aneurysms are associated with polycystic renal disease.

Ans:

True

[View All Answers](#)

Question - 69:

Berry aneurysms are associated with diabetes mellitus.

Ans:

False

[View All Answers](#)

Question - 70:

Berry aneurysms result from atheroma.

Ans:

False

[View All Answers](#)

Question - 71:

Berry aneurysm is most often found in the circle of Willis.

Ans:

True

[View All Answers](#)

Question - 72:

Abnormalities of blood clotting occur in von Willebrands disease.

Ans:

True

[View All Answers](#)

Question - 73:

Abnormalities of blood clotting occur in Christmas disease?

Ans:

True

[View All Answers](#)

Question - 74:

Abnormalities of blood clotting occur in sickle cell disease?

Ans:



False

[View All Answers](#)

Question - 75:

Abnormalities of blood clotting occur in thalassaemia.

Ans:

False

[View All Answers](#)

Question - 76:

Abnormalities of blood clotting occur in patients taking brufen.

Ans:

True

[View All Answers](#)

Question - 77:

Keratoacanthoma is histologically difficult to distinguish from squamous cell carcinoma.

Ans:

True

[View All Answers](#)

Question - 78:

In Keratoacanthoma, central necrosis is uncommon.

Ans:

False

[View All Answers](#)

Question - 79:

Keratoacanthoma is caused by a poxvirus.

Ans:

False

[View All Answers](#)

Question - 80:

Keratoacanthoma usually remits spontaneously without leaving scar.

Ans:

False

[View All Answers](#)

Question - 81:

Keratoacanthoma is also called molluscum sebaceum.

Ans:

True

[View All Answers](#)

Question - 82:

Keratoacanthoma causes increased risk of skin malignancy.

Ans:

False

[View All Answers](#)

Question - 83:

Squamous papilloma causes increased risk of skin malignancy.

Ans:

False

[View All Answers](#)

Question - 84:

Acanthosis nigricans causes increased risk of skin malignancy.

Ans:



False

[View All Answers](#)

Question - 85:

Solar keratosis causes increased risk of skin malignancy.

Ans:

True

[View All Answers](#)

Question - 86:

Bowens disease causes increased risk of skin malignancy.

Ans:

True

[View All Answers](#)

Question - 87:

In sickle cell disease, sickle cell crisis does not affect carrier of the disease.

Ans:

False

[View All Answers](#)

Question - 88:

In sickle cell disease, there is absence of HbA.

Ans:

True

[View All Answers](#)

Question - 89:

In sickle cell disease, the abnormality involves the alpha chain of the globin molecule.

Ans:

False

[View All Answers](#)

Question - 90:

In sickle cell disease, carrier has normal blood test.

Ans:

False

[View All Answers](#)

Question - 91:

In sickle cell disease, the disorder is a single gene disorder.

Ans:

True

[View All Answers](#)

Question - 92:

Keratoacanthoma predispose the squamous cell carcinoma.

Ans:

False

[View All Answers](#)

Question - 93:

Chronic ulceration predisposes the squamous cell carcinoma.

Ans:

True

[View All Answers](#)

Question - 94:

Chronic eczema predisposes the squamous cell carcinoma.

Ans:



False

[View All Answers](#)

Question - 95:

Bowens disease predisposes the squamous cell carcinoma.

Ans:

True

[View All Answers](#)

Question - 96:

Solar keratosis predisposes the squamous cell carcinoma.

Ans:

True

[View All Answers](#)

Question - 97:

Adenoma sebaceum seen in tuberous sclerosis is a type of hamartoma.

Ans:

True

[View All Answers](#)

Question - 98:

Hamartoma contains metaplastic cell types.

Ans:

False

[View All Answers](#)

Question - 99:

Hamartoma does not predispose to malignancy.

Ans:

True

[View All Answers](#)

Question - 100:

Hamartoma contains cells from all three germ layers.

Ans:

False

[View All Answers](#)

Question - 101:

Hamartoma is usually present at birth.

Ans:

True

[View All Answers](#)

Question - 102:

Amyloid causes renal failure.

Ans:

True

[View All Answers](#)

Question - 103:

Amyloid weakens the walls of blood vessels.

Ans:

True

[View All Answers](#)

Question - 104:

Amyloid is a basophilic substance, which can be stained with Congo red.

Ans:



False

[View All Answers](#)

Question - 105:

Amyloid occurs in medullary carcinoma of the thyroid.

Ans:

True

[View All Answers](#)

Question - 106:

Amyloid occurs in chronic inflammatory disorders.

Ans:

True

[View All Answers](#)

Question - 107:

Hyperplasia occurs in myocardium.

Ans:

False

[View All Answers](#)

Question - 108:

Hyperplasia occurs in retinal pigment epithelium.

Ans:

True

[View All Answers](#)

Question - 109:

Hyperplasia occurs in corneal endothelium.

Ans:

False

[View All Answers](#)

Question - 110:

Hyperplasia results in increased cell numbers.

Ans:

True

[View All Answers](#)

Question - 111:

Hyperplasia results in increased cell size.

Ans:

True

[View All Answers](#)

Question - 112:

In the presence of inflammation, complement proteins are raised.

Ans:

True

[View All Answers](#)

Question - 113:

In the presence of inflammation, fibrinogens are raised.

Ans:

True

[View All Answers](#)

Question - 114:

In the presence of inflammation, caeruloplasmin are raised.

Ans:



True

[View All Answers](#)

Question - 115:

In the presence of inflammation, ferritins are raised.

Ans:

True

[View All Answers](#)

Question - 116:

In the presence of inflammation, platelets are raised.

Ans:

True

[View All Answers](#)

Question - 117:

Multiple myeloma can give rise to metastatic calcification.

Ans:

True

[View All Answers](#)

Question - 118:

Pulmonary tuberculosis can give rise to metastatic calcification.

Ans:

False

[View All Answers](#)

Question - 119:

Renal failure can give rise to metastatic calcification.

Ans:

True

[View All Answers](#)

Question - 120:

Sarcoidosis can give rise to metastatic calcification.

Ans:

True

[View All Answers](#)

Question - 121:

Phthisical eye can give rise to metastatic calcification.

Ans:

False

[View All Answers](#)

Question - 122:

In cerebral infarction, the necrotic tissue is eventually replaced by Schwann cells.

Ans:

False

[View All Answers](#)

Question - 123:

In cerebral infarction, cortical blindness can result from infarction of the anterior cerebral artery.

Ans:

False

[View All Answers](#)

Question - 124:

In cerebral infarction, coagulative necrosis occurs in the brain tissue.

Ans:



False

[View All Answers](#)

Question - 125:

Cerebral infarction can result from thrombosis of the external carotid artery disease.

Ans:

False

[View All Answers](#)

Question - 126:

In cerebral infarction, the area of infarct tends to be wedge-shaped.

Ans:

True

[View All Answers](#)

Question - 127:

Factor V Leiden mutation may cause central retinal vein occlusion in a young patient.

Ans:

True

[View All Answers](#)

Question - 128:

Atrial fibrillation may cause central retinal vein occlusion in a young patient.

Ans:

False

[View All Answers](#)

Question - 129:

Antithrombin III deficiency may cause central retinal vein occlusion in a young patient.

Ans:

True

[View All Answers](#)

Question - 130:

Excess protein S may cause central retinal vein occlusion in a young patient.

Ans:

False

[View All Answers](#)

Question - 131:

Protein C deficiency may cause central retinal vein occlusion in a young patient.

Ans:

True

[View All Answers](#)

Question - 132:

In atherosclerosis, raised HDL is associated with atherosclerosis.

Ans:

False

[View All Answers](#)

Question - 133:

In atherosclerosis, proliferation of smooth muscle cells in the intima is typical.

Ans:

True

[View All Answers](#)

Question - 134:

In atherosclerosis, thinning of the intima is a feature.

Ans:



False

[View All Answers](#)

Question - 135:

In atherosclerosis, foamy macrophages are seen in type I plaque.

Ans:

False

[View All Answers](#)

Question - 136:

Atherosclerosis occurs in the retinal artery.

Ans:

False

[View All Answers](#)

Question - 137:

During wound healing, during collagen synthesis, lysine and proline are directly incorporated into the collagen molecules.

Ans:

False

[View All Answers](#)

Question - 138:

During wound healing, in infected wound collagen lysis is increase.

Ans:

True

[View All Answers](#)

Question - 139:

During wound healing, in scurvy the wound is weak due to increased activity of collagenase.

Ans:

False

[View All Answers](#)

Question - 140:

During wound healing, with absorbable suture, the wound strength decreases from the time of suturing.

Ans:

True

[View All Answers](#)

Question - 141:

During wound healing, the inflammation is greater for catgut than nylon

Ans:

True

[View All Answers](#)

Question - 142:

With regard to cell growth, achondroplasia does not affect membranous bone

Ans:

True

[View All Answers](#)

Question - 143:

With regard to cell growth, in metaplasia, there is a change of a type of differentiated cell to a type of undifferentiated cell

Ans:

False

[View All Answers](#)

Question - 144:

With regard to cell growth, the main stimulus for hypertrophy is hormonal

Ans:



False

[View All Answers](#)

Question - 145:

With regard to cell growth, hypocalcaemia can lead to enlargement of the parathyroid gland.

Ans:

True

[View All Answers](#)

Question - 146:

With regard to cell growth, ionizing radiation can cause atrophy.

Ans:

True

[View All Answers](#)

Question - 147:

The blood vessels surrounding an acutely inflamed site usually show progressive vasodilatation from the time of injury.

Ans:

False

[View All Answers](#)

Question - 148:

Autolytic changes in the nucleus are pathognomonic of necrosis.

Ans:

True

[View All Answers](#)

Question - 149:

Apoptosis does not usually cause inflammation.

Ans:

True

[View All Answers](#)

Question - 150:

Acute inflammation usually occurs around necrotic tissues.

Ans:

True

[View All Answers](#)

Question - 151:

Gangrene refers to tissue necrosis with or without infection.

Ans:

False

[View All Answers](#)

Question - 152:

The wound site becomes vascular as time passes.

Ans:

False

[View All Answers](#)

Question - 153:

Fibroblasts have a contractile function, which helps to produce a small scar.

Ans:

True

[View All Answers](#)

Question - 154:

The epidermal and dermal epithelia grow downwards along the suture track.

Ans:



True

[View All Answers](#)

Question - 155:

Monocytes clean the debris left by phagocytes.

Ans:

True

[View All Answers](#)

Question - 156:

Early suture removal does not affect the granulomatous response.

Ans:

False

[View All Answers](#)

Question - 157:

In ionizing radiation, undifferentiated tumors are usually more sensitive to radiation than differentiated tumors.

Ans:

True

[View All Answers](#)

Question - 158:

In ionizing radiation granulation tissue, formation is delayed in wounds, which have been radiated?

Ans:

True

[View All Answers](#)

Question - 159:

In ionizing radiation, cells experiencing hypoxia are vulnerable to radiation damage?

Ans:

False

[View All Answers](#)

Question - 160:

In ionizing radiation, mitosis does not occur in irradiated cells?

Ans:

False

[View All Answers](#)

Question - 161:

In ionizing radiation, cells in G1 phase of the cell cycle are most sensitive?

Ans:

False

[View All Answers](#)

Question - 162:

There are about 40 000 human genes and ESTs (expressed sequence tags) available on the Affymetrix Gene Chips for gene expression. Make some reasonable assumptions and estimate, for a normal organ, the number of genes that will appear to be expressed significantly greater than or less than the population norm (95% confidence interval).

Ans:

If we assume that each gene is independent and occurs in a normally distributed population, then one might expect approximately 5% of 40 000 or 2000 genes to appear over expressed or under expressed.

[View All Answers](#)

Question - 163:

Dr Jones is a pathologist who has recently developed a virtual microscope program for teenagers to use while they are hospitalized. He thinks that teens might use this program to learn more about their illness and the role that pathology plays in the hospital. He has heard that usability is important, so he asks a few of his colleagues to try the program out in their free time and to e-mail their comments to him. Describe 3 specific things that Dr Jones should have done differently in order; to more appropriately assess the usability of his program.

Ans:

- (1) Have teens test the program, not physicians;
- (2) select specific representative tasks for the assessment (not just "try the program out"); and



(3) observe the participants using the program and have them think aloud during the session rather than e-mail comments.

[View All Answers](#)

Question - 164:

To what 2 aspects of a usability assessment should the word "representative" apply?

Ans:

- (1) Representative tasks
- (2) Representative users

[View All Answers](#)

Question - 165:

What are the benefits of high-throughput expression analysis in molecular biological investigations?

Ans:

These techniques allow simultaneous analysis of the expression of many genes. Patterns of increased or decreased expression is associated with disease may then be identified and may contribute to improved diagnosis and prognosis.

[View All Answers](#)

Question - 166:

What is a "gene expression signature" for a tumor?

Ans:

A collection of genes that are expressed consistently higher or lower in the tumor than is the population norm for non-tumorous tissue of the same type.

[View All Answers](#)

Medical Most Popular & Related Interview Guides

- 1 : [Staff Nurse Interview Questions and Answers.](#)
- 2 : [Lab Technicians Interview Questions and Answers.](#)
- 3 : [Pharmaceutical Interview Questions and Answers.](#)
- 4 : [Pharmacist Interview Questions and Answers.](#)
- 5 : [Microbiology Interview Questions and Answers.](#)
- 6 : [Biotechnology Interview Questions and Answers.](#)
- 7 : [Healthcare Interview Questions and Answers.](#)
- 8 : [Nursing Interview Questions and Answers.](#)
- 9 : [Pharmacy Interview Questions and Answers.](#)
- 10 : [Physiotherapy Interview Questions and Answers.](#)

Follow us on FaceBook

www.facebook.com/InterviewQuestionsAnswers.Org

Follow us on Twitter

<https://twitter.com/InterviewQA>

For any inquiry please do not hesitate to contact us.

Interview Questions Answers.ORG Team

[https://InterviewQuestionsAnswers.ORG/
support@InterviewQuestionsAnswers.ORG](https://InterviewQuestionsAnswers.ORG/support@InterviewQuestionsAnswers.ORG)